## **MAGNESIUM CITRATE**

VALLEYCARE GASTROENTEROLOGY MEDICAL GROUP, INC.

Phillip Wolfe, MD

You are scheduled to have a colonoscopy on: \_\_\_\_\_

Located at: Hacienda Surgery Center You must arrive at \_\_\_\_\_\_A.M. 4626 Willow Rd, Suite 100, Pleasanton, CA 94588

Purchase TWO bottles of Magnesium Citrate, sold over the counter at local pharmacies. Purchase Dulcolax (you will need two tablets) to complete the colonoscopy pre/cleanse.

## THE DAY BEFORE YOUR COLONOSCOPY ~

## Please follow the instructions below from our office to prepare you for your procedure:

1. Start your clear liquid diet the whole day before your procedure. NO DAIRY PRODUCTS.

CLEAR LIQUID DIET: Water (stay hydrated)Tea or Coffee (without creamer)Clear juices without pulpFruit-flavored drinks (Kool-Aide, Crystal-Light, etc)Carbonated beverages/SodaSports drinks (Gatorade, All-Sport, PowerAde, etc)Broth/Bouillion/ConsommePlain/flavored gelatins (Jell-O)Fruit Ices/Italian IcesPopsicles (without fruit/fiber)

SAMPLE MENU:Breakfast: cup of broth, fruit juice, bowl of Jell-O, coffee or tea without creamer, water<br/>Lunch:<br/>tea or coffee, cup of broth, cup of fruit ice, fruit juice, water<br/>Snack:<br/>Popsicle, soda, Jell-O, water<br/>Dinner:<br/>Similar to lunch or breakfast, water

2. **Prep instruction for Magnesium Citrate prep. Follow the instructions below:** You must FINISH the entire solution. REMEMBER, if your colon is not well cleaned, the examination will not be useful and will need to be repeated.

The night before your procedure...

At 5:00 pm ....drink one bottle of Magnesium Citrate At 6:00 pm ....take 2 Dulcolax tablets At 8:00 pm ....drink the second bottle of Magnesium Citrate

Please be sure and drink 4 8-oz glasses of clear liquids or water after each bottle of Magnesium Citrate.

- 3. You may take medications with a few sips of water up to 2 hours before procedure.
- 4. If you are taking Coumadin, Plavix or any other blood thinning medications please discuss this with the gastroenterologist you are seeing in our office. All other medication should be taken as directed the morning of the procedure with very little water.
- 5. You will be receiving intravenous sedation or anesthesia for the Colonoscopy. YOU WILL NOT BE ABLE TO DRIVE THAT DAY. You must arrange for a responsible adult to take you home later that day.
- 6. Plan to be at the Out-Patient Surgery Center a total of 2-3 hours.

If you have any questions, please do not hesitate to call our office at 925-460-8167. www.valleycaregastro.com